

Analysis 3.1. Comparison 3 Active music therapy versus receptive music therapy, Outcome 1 Severity of depressive symptoms, clinician-reported (primary outcome; high=poor).	77
Analysis 3.2. Comparison 3 Active music therapy versus receptive music therapy, Outcome 2 Severity of depressive symptoms, patient-reported (primary outcome; high=poor).	77
Analysis 3.3. Comparison 3 Active music therapy versus receptive music therapy, Outcome 3 Quality of life (high=good).	78
Analysis 3.4. Comparison 3 Active music therapy versus receptive music therapy, Outcome 4 Leaving the study early.	78
APPENDICES	78
WHAT'S NEW	85
HISTORY	85
CONTRIBUTIONS OF AUTHORS	86
DECLARATIONS OF INTEREST	86
SOURCES OF SUPPORT	86
DIFFERENCES BETWEEN PROTOCOL AND REVIEW	87
INDEX TERMS	87

Music therapy for depression (Review) i

Copyright © 2017 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.



Trusted evidence.
Informed decisions.
Better health.

Cochrane Database of Systematic Reviews

[Intervention Review]

Music therapy for depression

Sonja Aalbers^{1,2}, Laura Fusar-Poli³, Ruth E Freeman⁴, Marinus Spreen⁵, Johannes CF Ket⁶, Annemiek C Vink⁷, Anna Maratos⁸, Mike Crawford⁹, Xi-Jing Chen¹⁰, Christian Gold¹¹

¹Social Work and Arts Therapies, University of Applied Sciences, Leeuwarden, Netherlands. ²Clinical, Neuro & Developmental Psychology, Vrije Universiteit Amsterdam, Amsterdam, Netherlands. ³Department of Brain and Behavioral Sciences, University of Pavia, Pavia, Italy. ⁴Psychiatry, Central and North West London NHS Foundation Trust, London, UK. ⁵School of Social Work and Art Therapies, Stenden University of Applied Sciences, Leeuwarden, Netherlands. ⁶Medical Library, Vrije Universiteit Amsterdam, Amsterdam.

10.1002/14651858.CD004517.pub3

[View Article Page](#)